



THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

### **Bioluminescence-based approach to monitor neural activity in freely moving *Drosophilalarvae***

**Citation for published version:**

Marescotti, M, Lagogiannis, K, Davies, W, Webb, B & Armstrong, JD 2016, 'Bioluminescence-based approach to monitor neural activity in freely moving *Drosophilalarvae*', 16th European Neurobiology of *Drosophila* Conference (Neurofly 2016), United Kingdom, 2/09/16 - 6/09/16.

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Publisher's PDF, also known as Version of record

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.





# brainwave-discovery

## Bioluminescence-based approach to monitor neural activity in freely moving *Drosophila* larvae

Marescotti M.<sup>1,2</sup>, Lagogiannis K.<sup>2</sup>, Davies W.<sup>1,2</sup>, Webb B.<sup>2</sup>, and Armstrong J.D.<sup>1,2</sup>  
 1.Brainwave-Discovery Ltd, Edinburgh, UK 2. University of Edinburgh

### Background:

Dissecting behavioural circuits requires us to examine activity in the brain as the animal processes sensory information and generates autonomous behaviour. With a few exceptions most current methods require a constrained preparation, which does not translate well to application in freely moving animals. We developed a non-invasive bioluminescence-based approach to address this problem; targeting the expression of the calcium-reporter Aequorin (AEQ) to the *Drosophila* larval nervous system to measure neural activity. We report conditions that significantly increase the sensitivity of this assay, allowing us to measure activity in Kenyon cells (100s) as well as in smaller populations of neurons, with clear signals obtained from less than 10 neurons in the living, unconstrained animal. Finally, we measure both spontaneous activity and evoked responses as the animal encounters specific environmental stimuli (e.g. tastes, odours).

### Methods: Bioluminescence principle






Figure 1

**Figure 1.** The principle of bioluminescence in *Drosophila* larvae in six steps. AEQ= Aequorin, CTZ= Coelenterazine, CLM= Coelenteramide. In red the covalent bound between CTZ and AEQ, that are broken when  $Ca^{++}$  bind AEQ.


### Results (I): Detection of neural activity in intact larvae



**Figure 2.** The set-up used to record bioluminescence from neurons in intact *Drosophila* larvae.




**Figure 3.** Detection of spontaneous activity of Kenyon cells from intact larvae. **a** and **c** represent the mean bioluminescence values, while **c**, **d-g** are representative individual examples.




**Figure 4.** Sensitivity range of the assay from 100s – 6 neurons. **a,b** and **i** show the mean bioluminescence values, **c-h** and **j** are representative individual examples.


### Results (II): Detection of neural activity from freely-crawling larvae



**Figure 5.** Simultaneous recording of neural activity in intact larvae and analysis of behavior using a photomultiplier in the UV range combined with IR video recording and tracking.



**Figure 6.** Pairing of neural activity in *Drosophila* larvae with movement in an arena.



**Figure 7.** Kenyon cells show increased activity in response to exposure to environmental  $CO_2$ . Bioluminescence from representative samples shown over time (before and after presenting  $CO_2$  to the larva).

### Conclusions and Ongoing work:

We developed a method allowing real-time analysis of neuronal activity in intact larvae while they are freely moving. Are aims for the future are. This system works for Kenyon cells and Dopaminergic neuron populations.

Ongoing work is looking at extending the range of stimuli in naïve and trained animals. We are also testing other neuronal populations.